

TB-500

Veterinary Use Only

Warning:

Check seal on vials and ampoules if seal is broken discard product.

Contents of box:

6 x 5ml Ampoules: Sterile saline solution
(Sodium Chloride Injection BP 0.9%)
6 x Vials: 10mg Thymosin Beta 4 Synthetic
Peptide

Sodium Chloride extraction:

Carefully open the top of the ampoule,
ensuring the contents are not spilled or
contaminated.

Place the assembled syringe in the dominant
hand and remove the protective cover, taking
care not to contaminate the needle.

Insert the needle into the ampoule and extract
2ml of sodium chloride solution.

Verify the correct dosage (2ml)

Dispose the remaining contents of the sodium
chloride solution responsibly

Important:

Prepare the vial for immediate use. If using
dosage that is less than the 2ml prepared (see
dog dosage) refrigerate the unused solution at
4 C

Preparation:

Remove the plastic protective cap of the sealed
vial containing the powder of peptides.

Insert the filled syringe containing the 2ml of
sodium chloride into the rubber top of the vial
and release the 2ml of sodium chloride into the
vial to produce 2ml of TB 500 solution. Remove
the syringe and gently rotate the vial until the
powder is completely dissolved. Ensure that the
solution is well mixed with all powder
thoroughly dissolved for maximum safety and
effectiveness.

Administration:

A Subcutaneous (S/C) injection.

Recommended needle size:

Horse 20g

Dog 23g

Recommended Dosage:

Horses

inject one 2ml vial each 7-10 days and repeat
with remaining 5 vials at 7-10 day intervals.

Dogs

inject 1ml of the 2ml vial prepared; refrigerate
the remaining unused 1ml solution at 4 C for
injection 7- 10 days later; repeat with each of
the remaining 5 vials 7-10 days later.

Adverse Reactions:

No adverse reactions have been reported to
date. Please report any adverse reactions
observed to: research@medivet.net.au

Contraindications:

No known contraindication exists with other
drugs, however the product has not been
thoroughly tested to know its full contraindica-
tions.

The product should not be used if the subject
animal is pregnant or breast feeding.

Storage: Store the unopened ampule and vials at
room temperature. Mixed vials must be
refrigerated at 4 C

MEDIVET



P.O. Box 3071 Bankstown Centro NSW 2200
Australia.

Phone: 1800 356 505

Fax: + 612 9708 0041

Email: research@medivet.net.au

Web: www.medivet.net.au